

Four Key Criteria for Genetic Material:

Information:	Contains instructions to build an organism.
Replication:	Capable of accurate copying (DNA replication).
Transmission:	Passed from parent to offspring and between cells during division.
Variation:	Accounts for differences within and between species.

Discovery of Genetic Material:

Early Hypotheses (Late 1800s):	August Weismann and Karl Nägeli proposed a biochemical basis for inheritance.
Chromosome Insight:	Chromosomes, composed of proteins and DNA, identified as carriers of genetic information.

Griffith's Bacterial Transformation Experiments:

Experiment Background:	☐ Studied <i>Streptococcus pneumoniae</i> . ☒ Type S (smooth, virulent) strains produce a polysaccharide capsule. ☒ Type R (rough, non-virulent) strains lack this capsule.
-------------------------------	--

Experimental Steps:

Step 1:	Injected live type R bacteria into a mouse → Mouse survived, no live bacteria found.
Step 2:	Injected live type S bacteria into a mouse → Mouse died, live type S bacteria found in blood.
Step 3:	Injected heat-killed type S bacteria into a mouse → Mouse survived, no live bacteria found.
Step 4:	Mixed heat-killed type S with live type R bacteria → Injected into a mouse → Mouse died, live type S bacteria found in blood.

Conclusion:

- ☐ Genetic material from heat-killed type S bacteria transformed live type R bacteria.
- ☐ This phenomenon was called "**transformation**" without knowing the biochemical nature of the transforming substance.

Transformation Concept:

Living type R bacteria transformed into type S, gaining the ability to produce a capsule.

This transformation indicated transfer of genetic material.

Avery, MacLeod, and McCarty

Focus: Investigated bacterial transformation, following up on Griffith's observations to identify the biochemical nature of the genetic material.

Experimental Approach:

Question: What substance from dead type S bacteria transforms live type R bacteria?

Purification Process: ☐ Purified macromolecules (proteins, DNA, RNA) from type S *Streptococcus pneumoniae*.
☐ Found only purified DNA could convert type R to type S bacteria initially.

Detailed Experiment:

Step 1:	Mixed purified DNA from type S bacteria with type R bacteria. Allowed DNA uptake by type R bacteria, converting some to type S.
Step 2:	Enzyme Treatments : ☐ DNase: Digests DNA. ☐ RNase: Digests RNA. ☐ Protease: Digests proteins.
Step 3:	Aggregated type R cells (non-transformed) removed by centrifugation.
Step 4:	Type S cells (transformed) remain in the supernatant.
Step 5:	Supernatant plated on growth media to observe bacterial colony formation.
Step 6:	Control plates (without DNA extract) showed no type S colonies.

Conclusion:

- ☐ DNA from type S bacteria alone could convert type R bacteria to type S, proving DNA as the genetic material.
- ☐ Elimination of transformation with DNase confirmed DNA's essential role.



By UmeshJagtap

Published 27th July, 2024.
 Last updated 27th July, 2024.
 Page 1 of 2.

Sponsored by **Readable.com**
 Measure your website readability!
<https://readable.com>

Hershey and Chase Experiment

Researchers: Alfred Hershey and Martha Chase (1952)

Objective: To determine whether DNA or protein is the genetic material in the T2 bacteriophage, a virus that infects *E. coli*.

☐ **Virus Structure Components:**

✓ **Capsid** (phage coat): Made entirely of protein, consisting of a head, sheath, tail fibers, and base plate.

✓ **DNA:** Found inside the head of the capsid.

Simplicity: Composed of only DNA and proteins.

Significance

Impact: The experiment provided convincing evidence that DNA, not protein, is the genetic material.

Scientific Legacy: This study was crucial in establishing DNA's role in heredity, greatly influencing molecular biology.

Experimental Design

Goal: To identify which component, DNA or protein, enters the bacterial cell and directs the synthesis of new viruses.

Key Insight: T2 phage injects its genetic material into the bacterial cell while the protein coat remains outside.

Methodology

Labeling: ☐ DNA labeled with ^{32}P (radioactive phosphorus).
☐ Protein labeled with ^{35}S (radioactive sulfur).

Infection Process: *E. coli* cells are infected with either ^{32}P -labeled phage or ^{35}S -labeled phage.

Shearing Force: Use a blender to detach phage coats from bacterial cells after allowing the phages to inject their genetic material.

Centrifugation: Separate heavier bacterial cells (pellet) from lighter phage coats (supernatant).

Detection: Measure the radioactivity in the pellet and supernatant using a Geiger counter.

Results

☐ ^{35}S (Protein): Majority found in the supernatant.

☐ ^{32}P (DNA): Majority found in the bacterial pellet.

Conclusion:

DNA enters the bacterial cell, not protein. This indicates that DNA is the genetic material responsible for the production of new viruses.



By UmeshJagtap

Published 27th July, 2024.

Last updated 27th July, 2024.

Page 2 of 2.

Sponsored by **Readable.com**

Measure your website readability!

<https://readable.com>